

Care at home Wound Formulary

July 2025

Wound care products should be ordered directly through the Igility Group Purchasing group.purchasing@igility.ca.

Note: If the product you need is not listed, please contact your wound consultant to discuss the possibility of an exception.

EXUDATE MANAGEMENT

Note: Avoid using foam dressings on feet or in conjunction with hydrogels.

Foam Dressings	<p>Lite Silicone Foam is appropriate for wounds with low levels of exudate. It may be used for up to 7 days or until exudate reaches within 0.5 cm of the dressing's edge.</p>		
	66800835	Allevyn Gentle Border Lite 10 cm x 10 cm	10/Box
	66800840	Allevyn Gentle Border Lite 15 cm x 15 cm	10/Box
	<p>Border Silicone Foam is suitable for wounds with moderate to heavy exudate and for managing skin tears. It can be used for up to 7 days or until exudate approaches within 0.5 cm of the dressing edge.</p>		
	66800269	Allevyn Gentle Border 7.5 cm x 7.5 cm	10/Box
	66800270	Allevyn Gentle Border 10 cm x 10 cm	10/Box
	66800272	Allevyn Gentle Border 12.5 cm x 12.5 cm	10/Box
	66800959	Allevyn Gentle Border 17.1 cm x 17.9 cm	10/Box
	<p>Border Silicone Foam is ideal for wounds with moderate to heavy exudate and for treating skin tears. The quadrilobed design allows the dressing to be used for up to 7 days or until exudate is visible in 3 of the 4 lobes.</p>		
	66801068	Allevyn Life 12.9 cm x 12.9 cm	10/Box
	66801069	Allevyn Life 15.4 cm x 15.4 cm	10/Box
	66801070	Allevyn Life 21 cm x 21 cm	10/Box
Alginate	<p>Post-Op Silicone Foam is designed for post-operative wounds with low to moderate exudate or when frequent dressing changes are needed. It can be used for up to 7 days or until exudate reaches within 0.5 cm of the dressing edge.</p>		
	66800900	Allevyn Gentle Border Post-op 10 cm x 20 cm	10/Box
	66800264	Allevyn Gentle Border Post-op 10 cm x 25 cm	10/Box
	<p>Non-Border Silicone Foam is suitable for wounds with moderate exudate and requires secondary fixation. It can be used for up to 7 days or until exudate approaches within 1.5 cm of the dressing edge.</p>		
	66802129	Allevyn Non-Border 10 cm x 10 cm	10/Box
	66802131	Allevyn Non-Border 15 cm x 15 cm	10/Box
	<p>Allevyn Classic is a non-silicone foam dressing designed for wounds with moderate to heavy exudate. It can remain in place for up to 7 days, depending on clinical conditions.</p>		
	66000044	Allevyn Adhesive Classic 12.5 cm x 12.5 cm	10/Box
	66000045	Allevyn Adhesive Classic 17.5 cm x 17.5 cm	10/Box
	<p>Calcium alginate is used to pack wounds and manage moderate to heavy exudate beneath a cover dressing. It may be left in place for up to 7 days or changed sooner if exudate accumulates in the wound or saturates the cover dressing.</p>		
	66000520	Algisite M 10 cm X 10 cm	10/Box
	66000522	Algisite M 2 cm x 30 cm	5/Box

Care at home Wound Formulary

July 2025

ANTIMICROBIAL

Antimicrobial dressings offer broad-spectrum activity and are highly effective in reducing microbial load.

Their use may be continued for 14 to 21 days to support wound improvement.

10% Povidone-Iodine Non-Adherent Sheet is appropriate for wounds with minimal exudate or those requiring drying. It may be used for up to 7 days or until the iodine color fades and must be secured with a cover dressing.

P01481	Iodine Dressing 5 cm x 5 cm	25/Box
P01512	Iodine Dressing 9.5 cm x 9.5 cm	25/Box

Cadexomer Iodine is indicated for sloughy, moist wounds and for managing biofilm. It remains effective for up to 3 days or until the dressing (e.g., Iodosorb) changes color to yellow or grey. A cover dressing is required.

66060630	Iodosorb Ointment 10 gm tube	4/Box
66001286	Iodosorb Powder 3 gm sachet	7/Box

Silver alginate is appropriate for wounds with moderate to heavy exudate.

It should not be used on dry wounds or those with minimal exudate.

The dressing may be used until the cover dressing becomes saturated or for a maximum of 7 days

CAD230	Silvercel Adherent Alginate Rope 2.5 cm x 30.5 cm rope	5/Box
CAD050	Silvercel Adherent Alginate 5cm x 5 cm	10/Box
CAD011	Silvercel Adherent Alginate 11 cm x 11 cm	5/Box

Nanocrystalline Silver Mesh can be used for all wounds, burns, and under NPWT. Use for 3-7 days.

66800545	Acticoat Flex 7 Rope 2.5 cm x 60 cm	5/Box
66800399	Acticoat Flex 3 10 cm x 10 cm (3-day wear)	12/Box
66800397	Acticoat Flex 7 12.5 cm x 12.5 cm (7-day wear)	5/Box

Nanocrystalline silver mesh is suitable for use on all wound types, including burns, and may also be applied under NPWT. It can be used for 3 to 7 days, depending on wound condition and exudate levels.

509341	UrgoTul Ag/Silver Contact Layer Dressing 10 cm x 12.5 cm	10/Box
--------	--	--------

Absorbent dressing with a fiber pad impregnated with silver sulfate, which forms a biocompatible gel upon contact with wound exudate. It is indicated for wound debridement and antimicrobial protection can be used up to 7 days or 1-3 days for heavy exudate or infection.

552155	UrgoClean AG Dressing 10 cm x 10 cm	10/Box
--------	-------------------------------------	--------

2% Polyhexamethylene Biguanide (PHMB) gauze dressing is indicated for wound packing, tunneling, and open surgical wounds. It should be used until the cover dressing becomes saturated, with a maximum wear time of 3 days.

Z7831AMD	AMD PHMB Packing Strips 0.63 cm x 0.91.4 m (1/4 inch)	10/Box
Z7832AMD	AMD PHMB Packing Strip 1.27 cm x 0.91.4 m (1/2 inch)	10/Box
Z7833AMD	AMD PHMB Packing Strip 2.51 cm x 0.91.4 m (1 inch)	10/Box
Z3332	AMD Kerlix Rolls 11.4 cm x 3.7 m	1 Roll

Methylene Blue/Gentian Violet aids in the removal of slough and eschar. **Pre-moisten the dressing for 5-10 minutes** before application. Use it for up to 3 days, and cover it with a dressing to maintain moisture.

HB2214	Hydrofera Blue Classic 5 cm x 5 cm	10/Box
HB4414	Hydrofera Blue Classic 10.2 cm x 10.2 cm	10/Box
HBT0906	Hydrofera Blue Classic 9 mm tunneling dressing	10/Box

Charcoal and silver dressing is designed for malodorous wounds and contains silver for antimicrobial action. Do not cut the dressing; instead, fold it to fit the wound. Use for up to 7 days, depending on the level of exudate

MAS105	Actisorb Ag 220 Charcoal 10.5cm x 10.5cm	10/Box
--------	--	--------

Care at home Wound Formulary

July 2025

Protective	Non-adherent covers help protect areas, keep them dry, and minimize the risk of sticking. Use for up to 7 days. Note: It does not promote moist wound healing.		
	7135	Primapore Adhesive Non-Woven Wound Dressing 8.3 cm x 6 cm	50/Box
	66000317	Primapore Adhesive Non-Woven Wound Dressing 10 cm x 8 cm	20/Box
Super-Absorbent	Super-Absorbent Dressing is used for absorbing high amounts of exudate. Use for up to 7 days. Either side of the dressing can be used. Ensure the dressing pad overlaps the edge of the wound. Do not cut the dressing.		
	PRD500-025	Kerramax Care Super-Absorbent Dressing 5 cm x 5 cm	10/Box
	PRD500-050	Kerramax Care Super-Absorbent Dressing 10 cm x 10 cm	10/Box
	PRD500-100	Kerramax Care Super-Absorbent Dressing 13.5 cm x 15.5 cm	10/Box
	PRD500-120	Kerramax Care Super-Absorbent Dressing 10 cm x 22 cm	10/Box
Skin Protectant	Skin Protectant Liquid Protective Film protects intact or damaged skin from wound exudate. Provides an effective barrier for up to 4 days when used alone or under non-adhesive dressings. Allow the liquid to dry completely before applying a cover dressing.		
	66800709	No Sting Barrier Film Spray 28 ml (single person use)	1 Bottle
	59420600	No Sting Barrier Film Wipe 1 ml	50/Box
	Adhesive Remover assists with the removal of adhesive dressings and tapes. Wipe off after use.		
	37436	Alkare wipe (adhesive remover)	50/Box
Antiseptic	Fabric Sheet with Slow-release Hydrogen Peroxide 0.3%, reduces bacterial growth Use: For moisture management and friction reduction between skin folds and other skin-to-skin contact areas, (e.g., finger/toe web spaces, hand contractures). Note: Additional 5cm is needed to allow the sheet to 'breathe' and dissipate the wicked moisture.		
	MSCWH219	Drigo-HP Wicking Sheet (10 x 144 roll)	1 Roll
Wound Cleanser	Chlorhexidine Gluconate (CHG) 2% Solution compounded with 70% alcohol is used to achieve and maintain dry, stable eschar (e.g., arterial wounds). Apply with friction for 30 seconds and reapply every 3 days. Do not apply to open wounds.		
	101.07	Soluprep antiseptic wipe – For smaller wounds	100/Box
	102.03	Soluprep antiseptic swab – For larger wounds	50/Box
	Betadine Povidone Iodine 10% is used on dry or open eschar to maintain an intact covering of a wound and eschar. Apply daily. Date the bottle when it is opened and discard the open bottle after 30 days.		
	1034105	Povidone Iodine (Betadine) 115 ml	1 Bottle
Wound Dressing	BIHOCL Pure Cleanse Wound Cleanser is a non-cytotoxic, antimicrobial solution designed for cleansing wounds, skin, and burns. Note: Rinsing is not required. Discard any unused solution 30 days after opening.		
	33055-118DC	BIHOCL Pure Cleanse Hypochlorous Acid Wound Cleanser 118ml	1 Bottle
	Normal Saline is used for wound cleansing. Hold the tip of the container 10-15 cm (4-6 inches) from the wound and squeeze the solution over the wound bed in a sweeping motion using all the solution. Discard the bottle after use.		
	T168000	100 ml Normal Saline- single use	25/Case

Care at home Wound Formulary

July 2025

Compression Wraps	<p>EdemaWear Tubular Compression Stocking delivers gentle longitudinal compression (approximately 10–20 mmHg). It is designed for continuous wear, day and night. Be sure to remove and reposition the stocking at least once daily. The stocking may remain in place for up to one week and should be washed in cold water and hung to dry. With proper care, it can be reused for up to six months.</p>		
	CA600001	Edema Wear Small up to 45 cm/ Navy (foot to knee). (2 stockings/ pack)	1 Package
<p>Compression Therapy: Non-elastic (short stretch) wraps may remain in place for up to 7 days, provided there is no slippage and wound exudate is effectively controlled.</p> <p>Note: Application should be performed only by healthcare professionals who have completed specialized training in compression therapy, in accordance with organizational policy.</p>			
<p>3M2094 Coban 2 Layer Short Stretch Compression 10 cm x 2.7 m 1/Box</p> <p>3M2794 Coban 2 Lite Layer Short Stretch Compression 10 cm wide 1/Box</p>			
Non-Adherent Contact Layer	<p>Wound Contact Layer: This petrolatum-impregnated dressing helps prevent wound bed trauma by minimizing adherence of the secondary dressing. It may remain in place for up to 7 days.</p> <p>Note: A cover dressing is required and must extend over the wound and at least 2 cm of the surrounding peri-wound skin.</p>		
	2015	Adaptic Non-Adhering Dressing 7.6 cm x 20.3 cm	24/ Box
Other	DUP59901	Dumex® Measure It® Wound Measurement Kit, Sterile	100/Box
	635-018-432	Medpro Cotton Tip Applicators	100/box
	195-7144301	Hypafix Conformable Adhesive Retention Tape 5 cm x 10m	1 Roll
	195-7144302	Hypafix Conformable Adhesive Retention Tape 10 cm x 10m	1 Roll
	7143616	Easifix Cohesive 8cm x 20 m	1 Roll